




Oxford University Hospitals
NHS Foundation Trust

Oramorph®

Information for patients



**Oxford Interstitial
Lung Disease Service**

Oramorph® 10mg/5ml Liquid is liquid morphine that is taken by mouth. Morphine is normally prescribed as a strong painkiller, but it can also work to help relieve your breathlessness.

For breathlessness, this medicine is usually prescribed in low doses of 2.5mg (1.25ml) to 5mg (2.5ml) and comes as a liquid for you to swallow. It is advisable to take it 15 to 20 minutes before any activity that makes you breathless, such as washing and dressing, or for the relief of shortness of breath after it has already developed.

Most people use Oramorph® 10mg/5ml Liquid 3 to 4 times a day, but it can be used every 4 hours if you need it, to control breathlessness symptoms.

What are the most common side effects?

Oramorph® 10mg/5ml Liquid may make you feel drowsy, but this is likely to improve as you continue with the treatment. Because of this, we advise you to take the first dose before going to sleep at night and to see how you feel in the morning. If you do not feel drowsy you can start to try using the Oramorph® 10mg/5ml Liquid during the day, when you feel you need it. If you do feel drowsy, then you should discuss this with your GP or Specialist ILD Nurse. You should also not drive when taking Oramorph® 10mg/5ml Liquid, until it is clear how drowsy it makes you feel.

Constipation is a very common side effect of taking Oramorph® 10mg/5ml Liquid. You may find that you need to take a laxative, which you can buy over the counter or discuss with your GP.

You may feel sick or vomit during the first 48 to 72 hours of starting this medication, but this usually improves after this time. If it continues, then you should seek advice from your GP or Specialist ILD Nurse.

Oramorph® 10mg/5ml Liquid can also cause itching of the skin. This may be helped by taking antihistamines, which you can buy over the counter or your GP can prescribe.

If you experience any of these side effects and they continue, despite taking the above steps, please let your GP or your Specialist ILD Nurse know.

You have been prescribed: mg in ml

An oral syringe will make it easier for you to measure your dose of Oramorph® 10mg/5ml Liquid. Please ask your local pharmacist to provide one for you.

Address Label

How to contact us

If you have any further questions please do get in touch. Useful phone numbers are listed below.

Oxford Interstitial Lung Disease Team (Churchill Hospital)

Lead Consultants' Secretary
Telephone: 01865 225 223

ILD Specialist Nurse
Email: oxford.ILD@ouh.nhs.uk

Further information

If you would like an interpreter, please speak to the department where you are being seen.

Please also tell them if you would like this information in another format, such as:

- Easy Read
- large print
- braille
- audio
- electronic
- another language.

We have tried to make the information in this leaflet meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They are happy to help.

Author: Oxford Interstitial Lung Disease Service
January 2024
Review: January 2027
Oxford University Hospitals NHS Foundation Trust
www.ouh.nhs.uk/information



Making a difference across our hospitals

charity@ouh.nhs.uk | 01865 743 444 | hospitalcharity.co.uk

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

