




Oxford University Hospitals  
NHS Foundation Trust

# Baker's Cyst

Information for patients



Oxford Anticoagulation  
and Thrombosis Service

## **Your ultrasound today has confirmed that you have a Baker's cyst.**

This is named after the British surgeon who first described the condition, William Marrant Baker. It is also known as a popliteal cyst.

## **What is a Baker's cyst?**

The term Baker's cyst is used to describe a fluid-filled swelling behind the knee. A Baker's cyst usually develops when the knee is damaged, which causes excess joint fluid to be produced. The excess fluid leads to an increase in pressure. This causes a bulge in the lining of the joint, out into the back of the knee, forming a Baker's cyst.

## **What causes it?**

Common conditions such as rheumatoid arthritis and osteoarthritis are often associated with a Baker's cyst.

Other types of knee injury, such as tears to the cartilage, may also be responsible. However, sometimes no cause can be found.

## **What are the symptoms?**

The size and amount of swelling produced by a Baker's cyst varies. A small cyst may give no symptoms; large cysts usually produce more problems.

### **Symptoms include:**

- discomfort or pain behind the knee
- tightness which increases when standing
- difficulty in bending the knee.

Occasionally the cyst will burst, causing pain and swelling behind the knee and down the calf muscle.

A Baker's cyst can sometimes be confused with a deep vein thrombosis (DVT) in the calf, but we can tell the difference by carrying out an ultrasound scan.

## **What treatment is available?**

Often the symptoms will get better without any treatment.

It is important to keep as mobile as possible. Elevating (raising) your leg may help to reduce the swelling. You can take a short course of over the counter anti-inflammatories to reduce pain and inflammation, but if a longer course is necessary this should be discussed with your GP. A support stocking may also help, which your GP can also advise you on.

If your symptoms do not improve it is important to see your GP.

If your symptoms are severe, a steroid injection into your knee may be recommended. You may also be referred to a rheumatologist or orthopaedic surgeon.

Surgery to remove the cyst is rarely necessary.

## **Where can I get further information?**

If you have any further questions, **please speak to your GP.**

## Further information

If you would like an interpreter, please speak to the department where you are being seen.

Please also tell them if you would like this information in another format, such as:

- Easy Read
- large print
- braille
- audio
- electronic
- another language.

We have tried to make the information in this leaflet meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They are happy to help.

Author: Thrombosis Team  
April 2022  
Review: April 2025  
Oxford University Hospitals NHS Foundation Trust  
[www.ouh.nhs.uk/information](http://www.ouh.nhs.uk/information)



*Making a difference across our hospitals*

[charity@ouh.nhs.uk](mailto:charity@ouh.nhs.uk) | 01865 743 444 | [hospitalcharity.co.uk](http://hospitalcharity.co.uk)

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

