




Oxford University Hospitals
NHS Foundation Trust

Video Assisted Anal Fistula Treatment (VAAFT)

Information for patients



This leaflet has been written to give you information about VAAFT (video assisted anal fistula treatment) surgery to treat an anal fistula.

What is an anal fistula?

An anal fistula is a small tunnel that develops between the end of the bowel and the skin of the buttock. It usually happens following an infection or abscess near the anus. You may be able to see or feel one or more holes near your back passage.

Anal fistulas can cause unpleasant symptoms, such as discomfort, skin irritation, and unpleasant discharge.

Anal fistulas can be simple or complex. Complex ones are deeper, going through the anal sphincter muscle, and may have extra passages or branches. The sphincter muscle is a ring around the back passage and gives control over the bowels. If it is damaged it weakens the control and can cause incontinence. Complex fistulas are difficult to treat because it is important not to damage this muscle. They can require several operations, over months or even years, to repair.

Surgery for anal fistula

It is not usually possible to know the full extent of the fistula before surgery and so decisions are often made whilst you are anaesthetised.

Simple or low fistulas

For simple fistulas the best treatment is to cut the fistula open by cutting through the skin directly. This usually involves cutting a small amount of the sphincter muscle, but the risk of incontinence is low. This creates a raw area that will heal by itself over a few weeks or months, depending on how long and deep the cut is.

Complex fistulas

Complex or deep fistulas cannot be treated like this because it would cause too much damage to the sphincter muscle and could result in loss of bowel control (bowel incontinence). A variety of other treatments are available, and your surgeon will discuss the options with you. The surgery may be planned in several stages over months or even years.

The first step is often to insert a Seton. A seton is a loose piece of silicon or surgical thread that is passed from the skin opening along the fistula, through the internal opening and out through the anus. It is then tied to form a loop that can stay in place for months or even years. Most people find a seton comfortable. You can go to the toilet and bathe or shower normally. Sometimes the outer part of the fistula, away from the muscle, is laid open at this time.

The seton allows the fistula to drain and helps it heal gradually from the inside. The seton usually controls the fistula but does not cure it. The seton is the first step of treatment; you may need several more procedures.

Further surgery

There are several options available for the further surgery, depending on the type of fistula and the preferences of the patient and surgeon. Combinations of treatments are often used.

Some options are:

- Remove the seton and hope the fistula closes or discharges a minimal amount. Sometimes the seton can be replaced with a very fine tube until the fistula closes.
- Core out the fistula track and close the internal opening using a section of the lining of the rectum (mucosal advancement flap).
- Close the fistula track with a biological plug, called an anal fistula plug.
- LIFT procedure (Ligation of Intersphincteric Fistula Tract). A cut is made near the back passage, and the space between the two rings of muscles is opened to reveal the fistula tract. This tract is then cut and tied off (ligated) on either side.
- Laser fistula ablation (FILAC). This is a minimally invasive technique. The fistula is cleaned, then a laser fibre is inserted into the fistula and laser energy is used to burn the inside of the fistula and seal it.
- VAAFT (Video Assisted Anal Fistula Treatment). A tiny camera is used to see inside the track. The fistula is cleaned and the lining of the fistula is burnt (cauterised) to help the fistula to shrink.

None of these methods are guaranteed to succeed at the first attempt, and sometimes multiple operations are required to eventually achieve healing of the fistula.

How does VAAFT work?

VAAFT is used:

- To see exactly where the fistula is going. It will show the internal opening and identify the track of fistula, including any branches. Because the surgeon uses direct vision, there is less risk of causing any damage.
- To lightly burn (cauterise) the inside of the fistula to encourage it to shrink.

VAAFT is most useful for complex fistulas with more than one tract or branch, and when the fistula is wide. It may need to be repeated more than once to encourage the fistula to shrink a bit each time.

What are the risks of VAAFT?

All procedures carry some risks. These will be explained to you before the operation. For VAAFT the risks include:

- Minor bleeding and pain are common. You may have a small amount of bleeding or ooze from the wound or anus after the operation. This is normal and nothing to worry about. It is a good idea to wear a small pad inside your pants for a week or two to protect your clothes.
- You might develop an infection with pus discharge from the external opening of the fistula. If this is associated with high temperature, you might need antibiotics to treat a wound infection.
- During VAAFT, fluid is used to wash the fistula tract. Occasionally this fluid escapes from the fistula and causes swelling in the buttock. This swelling usually goes down over 3-4 days and does not require any treatment.
- You may need further surgery to try to heal the fistula.

What to expect after your surgery

After your procedure you will be taken to the recovery ward where you will be looked after by specially trained nurses, under the direction of your anaesthetist. Once the effects of the general anaesthetic have worn off and you are comfortable, you will return to the ward.

Eating and drinking

You can eat and drink normally. We recommend a high fibre diet and drinking six to ten glasses of water every day. You will get additional information at the time of your discharge from hospital.

Moving around

You should start moving around as soon as possible after the procedure. This helps improve your recovery and reduces the risk of certain complications. Once back on the ward, you will be encouraged to get up and walk around.

Leaving hospital

You will be discharged from hospital on the same day.

Going to the toilet

From the day after your operation, you may be given laxatives to help soften the stools and keep you regular. There may be some discomfort and a little bleeding when you go to the toilet. This is to be expected. It is often a good idea to take pain killers 15 to 20 minutes before you try to open your bowels.

Looking after your wound at home

Before you go home your nurse will give you information about how to look after your wound at home. You will usually not need any dressings. There may be some fluid or discharge from the fistula. A small panty liner or pad will protect your underwear. You should take a bath or use a bidet, sitz bath, or shower nozzle to clean the area after opening your bowels.

How long should I stay off work and other normal activities?

- Most people need a week or two off work once they go home, but this depends on what you do and how large the fistula is.
- The time taken to get back to normal activities varies for different people and will depend on the surgery you have had.
- Do as much as you feel comfortable doing. If you need to take painkillers these may make you drowsy, so you should avoid driving or operating machinery.
- If lifting causes you discomfort, you should avoid it.
- You should not go swimming until your wound has healed as the chlorine in the water may affect wound healing and there is a chance of picking up or passing on an infection.
- You may find that vigorous exercise is uncomfortable. Start with gentle walking and build up your activity level gradually.
- You can have sex as soon as you feel comfortable to do so.
- You can drive as soon as you are comfortable enough to do an emergency stop safely. This is usually within a few days.

Further information

If you would like an interpreter, please speak to the department where you are being seen.

Please also tell them if you would like this information in another format, such as:

- Easy Read
- large print
- braille
- audio
- electronic
- another language.

We have tried to make the information in this leaflet meet your needs. If it does not meet your individual needs or situation, please speak to your healthcare team. They are happy to help.

Author: Miss Helen Jones
February 2025
Review: February 2028
Oxford University Hospitals NHS Foundation Trust
www.ouh.nhs.uk/information



Making a difference across our hospitals

charity@ouh.nhs.uk | 01865 743 444 | hospitalcharity.co.uk

OXFORD HOSPITALS CHARITY (REGISTERED CHARITY NUMBER 1175809)

